

LAWNWEED BREW

Kills weeds, not lawns

5-in-1! Controls Weeds, Disease, Moss, Algae, & Lichens in Turf

Works in cool weather (down to 50°F) Provides fast green-up of turf

Keep Out Of Reach Of Children
CAUTION (See Back Panel for Additional
Precaulionary Statements)



Derived from iron (Fe)



Concentrate

- ✓ Cure your weed, moss and disease* problems fast
- √Pre*- and Post-emergent activity of broadleaf weeds in one application
- √See results in hours
- √Rainfast in 3 hours
- √Can re-seed lawn 1 day after applying
- √Promotes healthy, weed- and disease*-free lawns

* Not for use in California

Manufactured under a license of W. Neudorff GmnHKG, Germany EPA Reg. No. 67702-26-4 EPA Est. No. 4-NY-1 Buyers Guarantee Limited to Label Claims.

©Bonide Products, LLC All Rights Reserved Distributed by: Bonide Products, LLC 6301 Sutliff Road, Oriskany, NY 13424 BN

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Sites: Use to selectively control/suppress broadleaf weeds, moss, algae*, lichens*, liverworts* and diseases* growing in lawns.

*Not for use in California

Pests:

Controls or suppresses weeds such as but not limited to:

Controls of suppresses weeds such as but not limited to.		
Black medic (Medicago lupulina) § Broadleaf plantain (Plantago major)* Bull thistle (Cirsium vulgare) Canada thistle (Cirsium arvense)* Common chickweed (Stellaria media) § ‡ Creeping buttercup (Ranunculus repens) Creeping Charlie/Ground	ivy (Glechoma hederacea)* • Dandelion (Taraxicum officinale) • Dovefoot geranium (Geranium molle)* • English daisy (Bellis perennis)§ • False dandelion (Hypochoeris radicata)* • Healall (Prunella vulgaris)* • Lawn burweed (Soliva ptero-sperma)*	

 Narrow-leaved plantain 	Slender
(Plantago lanceolata)*	speedwell(Veronica
• Oxalis (Oxalis spp.)*	filiformis)*)
Prostrate knotweed	White clover (<i>Trifolium</i>
(Polygonum aviculare)*	repens)
Shepherd's-purse	Wild chamomile (Matricaria)
(Capsella bursa-pastoris)*	chamomilla)*
Silverweed cinquefoil	Wild geranium (Geranium
(Potentilla anserina)*	robertianum)* `
 Speedwell spp.(including 	 moss, algae*, lichens*,
Persian Speedwell	liverworts*
(Veronica persica) and	

Controls or suppresses diseases such as but not limited to:

Dollar Spot*
 Rust*
 Snow Mold*

§ Only approved for suppression in California ‡Minimum application rate in California is 11.5 fl.oz./yd² or 9.8 ga/1000 sg.ft.

^{*}Not for use in California

APPLICATION DIRECTIONS

Shake well before using. Mix one part with 24 parts water (5 oz. in 1 gallon of water). Apply the mixed solution at a rate of 2.5-10 gallons/1000 ft² (2.8-11.5 fluid oz/yard²), or 5-10 gallons/1000 ft² (5.7-11.5 fl.oz./yd²) in California. Any standard hand-held or backpack sprayer can be used. Thoroughly spray weeds. Use a coarse nozzle setting to reduce drift. Uniform coverage is important. Repeat treatment in 3 to 4 weeks for best results.

§ Only approved for suppression in California

^{*} Not for use in California † Lower application rate not for use in California. Minimum application rate in California is 5.7 fl. oz per yard or 4.9 ga/1000 sq. ft.

To get the most complete kill of moss, or when moss is very thick, rake out as much moss as you can using a thatching rake or lawn rake before using this product.

To improve the appearance of lawn or turf, rake out dead moss a few days after treatment.

For Spot Treatment: Shake well before using. Mix one part with 24 parts water (5 fl. oz. in 1 gallon of water). Thoroughly spray weeds to point of runoff using any standard hand-held or backpack sprayer. Use a coarse nozzle setting to reduce drift. Repeat treatment in 3 to 4 weeks for best results.

For Pre-emergent control* of broadleaf weeds and crabgrass: Begin application when soil temperatures reach 50 to 55° F. Shake well before using. Mix one part with 24 parts water (5 fl. oz. in 1 gallon of water). Apply 5 to 10 gallons of spray solution per 1,000 sq. ft. depending on weed pressure. Reapply 3 to 6 weeks later for best results.

WHAT WILL I SEE?

Works quickly; dying weeds will be seen within hours of application. Treated weeds, moss, or algae* will turn brown or black. Occasionally a darkening of the grass leaf blades can occur after treatment; however, the grass will recover within a few days to weeks. May stain thatch (dead grass under lawn); this does not affect the health of the grass.

*Not for use in California

USE PRECAUTIONS

Do not apply to drought stressed grass; ensure lawn is well watered prior to application. Do not apply when the daytime temperature will exceed 85°F. Do not apply to bentgrass. If unsure of grass sensitivity, test spray a small area of your lawn. Do not apply to newly seeded areas prior to grass seed germination. Avoid spraying desirable plants. Any desirable plants or structures accidentally contacted should be rinsed with water. To avoid staining, wait until treated area dries before re-entry. Do not apply if rainfall is expected within 3 hours of application. Do not irrigate within 3 hours of application. Do not apply to the same area more than 4 time a year.

STORAGE and DISPOSAL

PESTICIDE STORAGE: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals. Store unused material, tightly closed, in original container only, away from open flame.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If nartly filled: Call your local solid waste agency, for

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

IF IN EYES: •Hold eye open and rinse

slowly and gently with water for 15-20 minutes.

•Remove contact lenses, if present, after the first 5 minutes then continue rinsing

FIRST AID

minutes then continue rinsing eye.

•Call a poison control center or doctor for treatment advice

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Call the poison control center at 1-800-222-1222.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes slight eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

LIMITED WARRANTY AND DISCLAIMER

Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for the purpose stated in the Direction for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. To the extent consistent with applicable law, this warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller. To the extent consistent with applicable law, the Seller shall not be liable for any such use. To the extent consistent with applicable law, the Seller makes no other warranties, either expressed or implied.