



# WEEDCLEAR™

## LAWN WEED KILLER

READY-TO-SPRAY

RESULTS IN HOURS

KILLS OVER

**250**  
WEEDS



DOLLARWEED DANDELION CLOVER CHICKWEED

**WON'T HARM LAWN GRASSES** WHEN USED AS DIRECTED

FOR RESIDENTIAL USE ONLY  
KEEP OUT OF REACH OF CHILDREN  
**CAUTION** See next panel for additional  
Precautionary Statements and First Aid.

NET CONTENTS 32 fl oz (1 quart) / 946 ml

<b>Active Ingredients:</b>	
2,4-D-chloroacetate ester	4.8%
Trifluralin	0.6%
Fluazifent	0.7%
Carbentrazone-methyl	0.1%
Inert Ingredients	93.8%
<b>Total</b>	<b>100.0%</b>



**This Product Contains:**  
0.228 lb 2,4-D-chloroacetate ester equivalent per gallon or 2.00%  
Inert Specific by AOC Method.



# WEEDCLEAR™

LAWN WEED KILLER  
READY-TO-SPRAY<sub>3</sub>

### PRODUCT FACTS

#### ⌚ Results in Hours

Kills major broadleaf lawn weeds including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge and many others.



### WHERE TO USE:



Ortho® WeedClear™ Lawn Weed Killer Ready-to-Spray<sub>3</sub> is intended for use only on residential lawns including Kentucky bluegrass, perennial ryegrass, fescues, common bermudagrass, hybrid bermudagrass, bahiagrass, bentgrass, buffalograss, centipedegrass, zoysiagrass and St. Augustinegrass (except 'Floratum').

Do not allow people or pets to enter the treated area until sprays have dried.

Questions & Comments  
Call 1-800-225-2883  
or [ortho.com](http://ortho.com)

Stop! Read the entire label first. Observe all precautions and follow directions carefully.

FPO

0 71549 04491 6  
0505



If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

©2019 The ORTHO Group. World rights reserved.  
Sold by The ORTHO Group,  
PO Box 190, Marysville, OH 43040  
EPA Reg. No. 2217-910-239  
EPA Est. No. 239-1A-3



LB13881

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

<b>PESTICIDE STORAGE</b>	Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.
--------------------------	--

<b>PESTICIDE DISPOSAL AND CONTAINER HANDLING</b>	Nonrefillable container. Do not reuse or refill this container. <b>If empty:</b> Place in trash or offer for recycling if available. <b>If partly filled:</b> Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.
--	---

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. When using this product, wear long-sleeved shirt and long pants, socks, shoes, and gloves.

### Personal Hygiene Statements

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

## FIRST AID

<b>IF ON SKIN</b>	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice.
-------------------	---

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-465-5161 for emergency medical treatment advice.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## PHYSICAL AND CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.



If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

©2019 The ORTHO Group. World rights reserved.  
Sold by **The ORTHO Group**  
P.O. Box 190, Marysville, OH 43040  
EPA Reg. No. 2217-910-239  
EPA Est. No. 239-IA-3

 **Questions & Comments**  
Call 1-800-225-2883  
or [ortho.com](http://ortho.com) 

LB13881

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### PRECAUTIONS AND RESTRICTIONS :

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.
- Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans.
- Do not apply when windy
- To prevent product run-off, do not over water the treated areas to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.

## WHERE TO USE

Ortho® WeedClear™ Lawn Weed Killer Ready-To-Spray, is intended for use only on residential (home) lawns including Kentucky bluegrass, perennial ryegrass, fescues, common bermudagrass, hybrid bermudagrass, bahiagrass, bentgrass, buffalograss, centipedegrass, zoysiagrass and St. Augustinegrass (except 'Floritam').

## AMOUNT TO USE

Use the amount indicated for your type of lawn:

**For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (except 'Floritam'):** This container covers approximately 10,600 square feet.

**For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass:** This container covers approximately 5,300 square feet.

## HOW TO USE THE READY-SPRAY® SYSTEM

1. Measure the total area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet.
2. See below directions for the ready-to-use hose-end sprayer.
3. Spray evenly over the entire measured area.

<b>CONNECT</b>	<ul style="list-style-type: none"><li>• Shake well before using.</li><li>• Connect sprayer to hose.</li><li>• Turn on water.</li><li>• Extend hose to farthest area of the measured area to be treated.</li></ul>
<b>SPRAY</b>	<ul style="list-style-type: none"><li>• To begin spraying point nozzle in the direction you want to spray.</li><li>• Slide thumb activation switch forward to ON position.</li><li>• Spray evenly over measured area.</li></ul>
<b>FINISH</b>	<ul style="list-style-type: none"><li>• To STOP spraying, slide thumb activation switch backward to OFF position.</li><li>• Turn off water.</li><li>• Relieve water pressure by sliding thumb switch to ON position until water slows to a drip. Then, slide switch back to OFF position.</li><li>• Disconnect sprayer from hose.</li></ul>

## WHEN TO USE

- Spray when broadleaf weeds are young and actively growing for best results.
- Ortho® WeedClear™ Lawn Weed Killer Ready-To-Spray, may be applied in the spring, summer or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.
- Use broadcast (entire areas) applications or follow-up applications at 30 day intervals for more mature weeds, for dense infestations, and for adverse environmental conditions.
- Spot treatments during the summer may be appropriate for sparse infestations, as a follow-up treatment, or any time broadleaf weeds are susceptible.
- **For Established Lawns:** Apply when daytime temperatures are below 90°F.
- **For Newly Seeded Areas:** Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- **For Newly Sodded, Sprigged or Plugged areas:** Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
- **Re seeding:** Treated areas can be reseeded at 2 weeks after the application of this product.
- **Limitations:** The maximum rate is 6.0 fl.oz. of product per 1,000 sq.ft. per application, and the maximum number of broadcast applications is two (2) per year with a minimum of 30 days between applications. The maximum seasonal rate is 12.0 fl.oz. of product per 1,000 sq.ft. excluding spot treatments.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated areas to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.
- **Irrigation:** Adequate soil moisture and favorable growing conditions enhance the performance of this product. If dry conditions exist, schedule irrigations (watering) before and after the application. Avoid applications of this product when turf is under stress since injury may result.
- **Mowing:** Avoid mowing within 1 to 2 days before and after the application of this product.
- **Non-target plants:** Do not apply this product to carpetgrass, dichondra, food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.

## BROADLEAF WEEDS CONTROLLED

Ortho® WeedClear™ Lawn Weed Killer Ready-To-Spray, will control or suppress the following broadleaf weeds.

Amaranth, purple; Amaranth, slender; Aster, heath; Aster, white heath; Aster, purple; Aster, white prairie; Artichoke; Austrian fieldcress; Bedstraw, catch-weed; Bedstraw, smooth; Bedstraw, northern; Bedstraw, pineywoods; Beggar-ticks, hairy; Betony, Florida; Bindweed, field; Bindweed, hedge; Bitter Wintercress; Bitter-cress, hairy; Bitterweed; Black-eyed Susan; Black medic; Blood-flower milkweed; Blue lettuce; Brassbuttons; Bristly ox-tongue; Broomweed; Buckhorn; Bull-nettle; Bur-clover; Burdock; Burweed, lawn; Buttercup, bulbous; Buttercup, creeping; Buttonweed, common (poorjoe); Buttonweed, Virginia; Carolina geranium; Carpetweed; Catnip; Catsear, spotted (false dandelion); Chamber bitter; Chickweed, common; Chickweed, mouse-ear; Chickweed, sticky; Chicory; Cinquefoil; Clover, crimson; Clover, Alyce; Clover, hop; Clover, rabbit-foot; Clover, red; Clover, strawberry; Clover, yellow sweet; Clover, white sweet; Clover, white; Cockle; Cocklebur; Cocklebur, spiny; Common mullein; Corn spurry; Creeping beggar-weed; Creeping Jenny (creeping Charlie); Cupid's shaving brush; Cudweed; Daisy, English; Daisy, oxeye; Dandelion, common; Dandelion, false; Day-flower; Deadnettle; Dichondra; Dock, broadleaf; Dock, curly; Dogbane; Dog-fennel; Dollarweed (Pennycort); Dove-weed; Elderberry; Falseflax; False sunflower; Fiddleneck; Filaree, redstem; Filaree, whitestem; Fleabane, daisy (annual); Fleabane, rough; Galinsoga, hairy; Galinsoga, small flower;

(Continued) 3

### BROADLEAF WEEDS CONTROLLED (Continued)

Goldenrod; Ground ivy; Groundsel; Gumweed; Hairy fleabane; Hawkweed, meadow; Hawkweed, mouse-ear; Hawkweed, orange; Hawkweed, yellow; Heaball; Heart-leaf drymary; Hemp; Henbit; Hoary-cress; Horsenettle; Horseweed; Jimsonweed; Johnny-jumpup violet; Knawel; Knotweed, prostrate; Kochia; Lambsquarters; Lespedeza; Mallow, alkali; Mallow, bristly; Mallow, common; Mallow, dwarf; Mallow, Venice; Marcella; Matchweed; Mexican-weed; Moneywort; Morningglory, bigroot; Morningglory, ivyleaf; Morningglory, red; Morningglory, tall; Morningglory, woolly; Mugwort; Mustard, ball; Mustard, black; Mustard, blue; Mustard, hedge; Mustard, Indian; Mustard, tansy; Mustard, tumble; Mustard, white; Mustard, wormseed; Nettle, stinging; Nettle, tall; Nettle, wood; Parsley-piert; Parsnip; Pearlwort; Pennygrass; Pepperweed, perennial; Pepperweed, Virginia (Peppergrass); Pigweed, prostrate; Pigweed, redroot; Pigweed, Russian; Pigweed, smooth; Pigweed, tumble; Pineapple weed; Plains coreopsis (tickseed); Plantain, black-seed; Plantain, bracted; Plantain, broadleaf; Plantain, buckhorn; Plantain, hoary; Plantain, narrow-leaf; Plantain, slender; Plantain, woolly; Poison ivy; Poison oak; Pokeweed; Prairie sunflower; Prickly lettuce (compass plant); Prickly sida; Prostrate knotweed; Puncturevine; Purslane, common; Pusley, Brazilian; Pusley, Florida; Pusley, large flower; Ragweed, bur; Ragweed, common; Ragweed, lance-leaf; Ragweed, western; Redstem filaree; Rough cinquefoil; Scarlet pimpernel; Shepherd's purse; Smartweed, ladythumb; Smartweed, pale; Smartweed, Pennsylvania; Smooth chaff-flower; Smooth dock; Sorrel, red (sheep); Sowthistle, annual; Sowthistle, spiny; Spanish needles; Speedwell, bi-lobed; Speedwell, common; Speedwell, corn; Speedwell, ivyleaf; Speedwell, Persian; Speedwell, purslane; Speedwell, slender; Speedwell, snow; Speedwell, thyme-leaf; Spiny amaranth; Spurge, spotted; Spurge, prostrate; Star of Bethlehem; Strawberry, Indian mock; Tansy ragwort; Tanweed; Thistle, blessed; Thistle, bull; Thistle, Canada; Thistle, Hodman's; Thistle, musk; Thistle, plumeless; Thistle, prairie; Thistle, Russian; Thistle, Scotch; Thistle, wavy-leaf; Thistle, yellow; Trailing crown-vetch; Velvetleaf; Vervain, blue; Vervain, hoary; Vervain, prostrate; Vervain, tall; Vetch, bird; Vetch, hairy; Vetch, milk; Vetch, narrow-leaf; Vetch, wild; Virginia creeper; Western clematis; Western salsify; Wild aster; Wild buckwheat; Wild carrot; Wild four-o'clock; Wild garlic; Wild lettuce; Wild marigold; Wild mustard; Wild onion; Wild parsnip; Wild radish; Wild rape; Wild strawberry; Wild sweet potato; Wild violet; Woodsorrel, creeping; Woodsorrel, violet; Woodsorrel, yellow (*Oxalis*); Woolly croton; Wormseed; Yarrow, common; Yarrow, western; Yarrow, yellow; Yellow Rocket

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

<b>PESTICIDE STORAGE</b>	Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.
<b>PESTICIDE DISPOSAL AND CONTAINER HANDLING</b>	Nonrefillable container. Do not reuse or refill this container. <b>If empty:</b> Place in trash or offer for recycling if available. <b>If partly filled:</b> Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. When using this product, wear long-sleeved shirt and long pants, socks, shoes, and gloves.

#### Personal Hygiene Statements:

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

## FIRST AID

<b>IF ON SKIN</b>	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice.
-------------------	---

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-465-5161 for emergency medical treatment advice.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## PHYSICAL AND CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

## MODO DE EMPLEO

Es una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta.

### PRECAUCIONES Y RESTRICCIONES

- No aplique este producto de forma que entre en contacto con otras personas o mascotas, ya sea directamente o al ser arrastrado por el viento. Durante la aplicación, mantenga alejadas del área a las personas y las mascotas.
- No permita que las personas ni las mascotas ingresen al área tratada hasta que el rociado se haya secado.
- No aplique como una niebla fina por los posibles daños a las plantas que no desea eliminar.
- No aplique directamente ni cerca de agua, drenajes pluviales, cunetas, alcantarillas o zanjas de desagüe.
- No aplique a menos de 25 pies (8 m) de distancia de ríos, estanques de peces, lagos, arroyos, pantanos, marismas, estuarios, bahías y océanos.
- No aplique cuando haga viento.
- Para evitar escurrimientos del producto, no riegue las áreas tratadas al grado que escurra ni lo aplique cuando esté lloviendo o si se pronostican lluvias para ese día. Enjuague el aplicador sobre el área del césped o el jardín, únicamente.

## DÓNDE USARLO

El Herbicida para césped WeedClear™ de Ortho® listo para rociar, está indicado para su uso exclusivo en céspedes residenciales (domésticos), como azul de Kentucky, raigrás perenne, cañueñas o festucas, bermuda común, bermuda híbrido, bahía, agrostis, búfalo, ciempiés, zoysia y San Agustín (excepto la variedad 'Floratum').

## CANTIDAD A UTILIZAR

Use la cantidad indicada para su tipo de césped:

**Para céspedes agrostis, bermuda híbrido, búfalo, ciempiés y San Agustín (excepto la variedad Floratam):** Con el contenido de este envase se cubren aproximadamente 10,600 pies cuadrados (985 m<sup>2</sup>).

**Para céspedes Azul de Kentucky, bahía, bermuda común, cañuelas o festucas, raigrás perenne y zoysia:** Con el contenido de este envase se cubren aproximadamente 5,300 pies cuadrados (992 m<sup>2</sup>).

## CÓMO USAR EL SISTEMA READY-SPRAY®

1. Mida el área total a rociar. Para áreas rectangulares, multiplique el largo por el ancho para obtener el área total.
2. Consulte las instrucciones del rociador para manguera listo para usar a continuación.
3. Rocíe de manera uniforme sobre toda el área medida.

<b>CONECTE</b>	<ul style="list-style-type: none"><li>• Agite bien el producto antes de usarlo.</li><li>• Conecte el rociador a la manguera.</li><li>• Abra la llave del agua.</li><li>• Extienda la manguera hasta el extremo más alejado del área medida a tratar.</li></ul>
<b>ROCÍE</b>	<ul style="list-style-type: none"><li>• Para comenzar, apunte la boquilla en la dirección que desee rociar.</li><li>• Deslice el interruptor de activación hacia adelante con el pulgar hasta la posición de ON (abierto).</li><li>• Rocíe de manera uniforme sobre el área medida.</li></ul>
<b>TERMINE</b>	<ul style="list-style-type: none"><li>• Para DEJAR de rociar, deslice el interruptor de activación hacia atrás con el pulgar hasta la posición de OFF (cerrado).</li><li>• Cierre la llave del agua.</li><li>• Libere la presión del agua deslizando el interruptor con el pulgar hasta la posición de ON (abierto), hasta que el chorro del agua se reduzca a un goteo.</li><li>• A continuación, deslice de nuevo el interruptor hasta la posición de OFF (cerrado).</li><li>• Desconecte el rociador de la manguera.</li></ul>

## CUÁNDO USARLO

- Para obtener los mejores resultados, rocíe cuando las malezas de hoja ancha sean pequeñas y estén en crecimiento activo.
- El Herbicida para césped WeedClear™ de Ortho® listo para rociar, puede aplicarse en primavera, verano o otoño. Los tratamientos en primavera y otoño, con condiciones de humedad de la tierra adecuadas, son preferibles a los tratamientos de verano. En general, las aplicaciones por voleo en malezas más viejas y estresadas por la sequía durante el verano son menos eficaces.
- Recorra a aplicaciones por voleo (áreas enteras) o de seguimiento a intervalos de 30 días para tratar las malezas más maduras y las infestaciones densas, y cuando imperen condiciones ambientales adversas.

- Los tratamientos focalizados durante el verano pueden ser apropiados para las infestaciones dispersas como un tratamiento de seguimiento o en cualquier momento en que las malezas de hoja ancha sean vulnerables.
- **Para céspedes establecidos:** Aplique cuando la temperatura esté por debajo de los 90 °F (32 °C).
- **Para áreas recién sembradas:** Aplique este producto a los céspedes recién sembrados cuando estén bien establecidos (aproximadamente 6 semanas después del nacimiento de las plántulas) o después del tercer corte.
- **Para áreas recién sembradas, plantadas o cubiertas con césped:** Aplique este producto 3 o 4 semanas después de sembrar, plantar o cubrir con césped.
- **Resembrado:** Las áreas tratadas pueden volverse a sembrar 2 semanas después de la aplicación de este producto.
- **Limitaciones:** La proporción máxima es de 6 onzas líquidas (177 ml) de producto por cada 1,000 pies<sup>2</sup> (93 m<sup>2</sup>), por aplicación, y el número máximo de aplicaciones por voleo es de dos (2) al año, con un mínimo de 30 días entre una aplicación y otra. La proporción máxima por temporada es de 12 onzas líquidas (356 ml) de producto por cada 1,000 pies<sup>2</sup> (93 m<sup>2</sup>), excluyendo los tratamientos focalizados.
- No aplique como una niebla fina por los posibles daños a las plantas que no desea eliminar.
- No use este producto en plantas que no desee eliminar ni cerca de ellas, incluyendo el contacto del riego con los sistemas de raíces expuestos o los brotes fortuitos dentro de la línea de goteo de árboles y arbustos deseables, pues pueden sufrir daños.
- No aplique directamente ni cerca de agua, drenajes pluviales, cunetas, alcantarillas o zanjas de desagüe. No aplique a menos de 25 pies (8 m) de distancia de ríos, estanques de peces, lagos, arroyos, pantanos, marismas, estuarios, bahías y océanos. No aplique cuando haga viento. Para evitar escurrimientos del producto, no riegue el/ las área(s) tratada(s) al grado que escorra ni lo aplique cuando esté lloviendo o si se pronostican lluvias para ese día. Enjuague el aplicador sobre el área del césped, únicamente.
- **Riego:** Una humedad de la tierra adecuada y condiciones de crecimiento favorables mejoran el rendimiento de este producto. De imperar condiciones secas, programe una irrigación (el riego) para antes y después de la aplicación. Evite aplicar este producto cuando el césped esté estresado, pues puede sufrir daños.
- **Corte:** No corte el césped 1 a 2 días antes o después de la aplicación de este producto.
- **Plantas que no pretende eliminar:** No aplique este producto a céspedes de las variedades alfombra o dicandra, a cultivos comestibles (huertos, plantas frutales y hortalizas), forrajes o plantas ornamentales, incluyendo flores, árboles, arbustos, setos, plantas leñosas ornamentales, cubiertas vegetales establecidas en jardines, y otras plantas que no se desee eliminar.

## ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua, la comida ni el pienso para animales al almacenar o eliminar el producto.	
<b>ALMACENAMIENTO DEL PESTICIDA:</b>	Guarde el producto en su envase original, en un lugar de almacenamiento bajo llave e inaccesible para los niños o las mascotas. Proteja del congelamiento.
<b>ELIMINACIÓN DEL PESTICIDA Y MANEJO DEL ENVASE:</b>	Este envase no se puede rellenar. No vuelva a usar ni a llenar este envase. <b>SI ESTÁ VACÍO:</b> Tirelo a la basura o, si está disponible, ofrézcalo para reciclaje. <b>SI ESTÁ PARCIALMENTE LLENO:</b> Llame a su agencia de desechos sólidos local para obtener instrucciones sobre cómo eliminar el producto. Nunca elimine el producto no utilizado en drenajes internos o externos.

## AVISOS DE PRECAUCIÓN

### RIESGOS PARA LOS SERES HUMANOS Y LOS ANIMALES DOMÉSTICOS

**PRECAUCIÓN:** Es nocivo si se absorbe a través de la piel. Evite el contacto con la piel, los ojos o la ropa. Vista una camisa de manga larga, pantalones largos, medias o calcetines, zapatos y guantes al usar este producto.

#### **Dedaciones de higiene personal:**

Lávese bien con agua y jabón después de manipular el producto, y antes de comer, beber, masticar chicle o consumir tabaco. Quítese y lave la ropa contaminada antes de volver a usarla.

## PRIMEROS AUXILIOS

### SI CAE EN LA PIEL:

Quítese la ropa contaminada.  
Enjuague la piel de inmediato con abundante agua por 15 a 20 minutos.  
Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

Tenga el envase o la etiqueta del producto con usted cuando llame al centro de control de envenenamientos o al médico, o al acudir a recibir tratamiento. También puede llamar al 1-877-465-5161 para obtener información sobre tratamiento médico de emergencia.

## RIESGOS MEDIOAMBIENTALES

Este pesticida es tóxico para los peces y los invertebrados acuáticos, y puede afectar de manera adversa a las plantas que no se pretende eliminar. Para proteger el medio ambiente, no permita que el pesticida penetre o se escurra en drenajes pluviales, zanjas de desagüe, cunetas o aguas superficiales. La aplicación de este producto en un clima tranquilo, cuando no se pronostiquen lluvias para las próximas 24 horas, asegurará que ni el viento ni la lluvia arrastren o laven el pesticida fuera del área de tratamiento. Enjuagar el equipo usado para la aplicación sobre el área tratada ayudará a evitar escurrimientos hacia cuerpos de agua o sistemas de drenaje.

Este producto químico tiene propiedades y características que se relacionan con sustancias químicas detectadas en aguas del subsuelo. El uso de este producto químico en áreas donde la tierra es permeable, en particular donde el manto freático está cerca de la superficie, puede acarrear la contaminación del agua del subsuelo. La aplicación alrededor de una cisterna o pozo puede ocasionar la contaminación del agua para beber o de la del subsuelo.

## RIESGOS FÍSICOS Y QUÍMICOS

Combustible. No utilice ni almacene el producto cerca de una fuente de calor o llama abierta.

**ADVERTENCIA:** El comprador asume todos los riesgos derivados del uso, el almacenamiento o el manejo de este producto si no se siguen las instrucciones de la etiqueta.