

10921 Weed-Out Plus Lawn Fertilizer

ferti·lome®



# Weed-Out Plus Lawn Fertilizer

## 25-0-4

### Contains Trimec® Herbicide

Controls dandelion, chickweed, knotweed, plantain, henbit, spurge, and many other broadleaf weeds which are listed on this label in St. Augustine, centipedegrass, bahiagrass, bermudagrass, zoysiagrass, bluegrass, perennial ryegrass, fescue and bentgrass lawns.

**ACTIVE INGREDIENTS**

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid.....	0.29%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	0.17%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid) .....	0.07%
INERT INGREDIENTS.....	99.47%
TOTAL.....	100.00%

**THIS PRODUCT CONTAINS:**

4.80 lbs. or 0.24% (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.  
2.80 lbs. or 0.14% 2,4-dichlorophenoxyacetic acid equivalent per ton.  
1.20 lbs. or 0.06% 3,6-dichloro-o-anisic acid equivalent per ton.  
Isomer Specific by AOAC Methods.  
TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

**KEEP OUT OF REACH OF CHILDREN**  
**SEE BACK PANEL FOR FIRST AID, PRECAUTIONARY STATEMENTS**  
**AND DIRECTIONS FOR USE**

**CAUTION**

**NET WEIGHT 20 LBS. (9.8 KG)**

**Weed-Out**  
**Plus Lawn Fertilizer**

**Weed-Out Plus Lawn Fertilizer**

**Weed-Out Plus Lawn Fertilizer**

**Covers Up To**  
**5,000 Sq. Ft.**



7 32221 10921 6



# Weed-Out Plus Lawn Fertilizer

**STOP!** READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt, long pants, shoes plus socks and chemical-resistant gloves (made of any waterproof material). User should wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet.

### FIRST AID

<b>If In Eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a Poison Control Center or doctor for treatment advice.</li> </ul>
<b>If On Skin Or Clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>Call a Poison Control Center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a Poison Control Center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>

Have the product container or label with you when you calling a Poison Control Center or doctor or going for treatment.

In the event of a medical or chemical emergency contact ChemTel  
North America 1-800-255-3924

You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### General Precautions and Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application
- Do not allow people or pets to enter the treated area until dusts have settled.

### USE PRECAUTIONS

Do not exceed specified dosages for any area. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

#### Other Use Precautions:

- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

### INSTRUCTIONS

**ferti•lome® WEED-OUT® PLUS LAWN FERTILIZER** controls a wide range of lawn weeds, including Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many others commonly found in home lawns. In addition, a 25-0-4 fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Do not use on Dichondra. Do not over apply. Under conditions of adequate moisture, high temperatures and high humidity, over-application may cause minor transitory discoloration on St. Augustinegrass. Do not use this product on "Floratam" St. Augustinegrass.

- Mow lawn to normal height 1 to 2 days before application.
- Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (See Step 4).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto non-target plants.
- Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

### HOW MUCH TO APPLY

**For Kentucky Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Zoysiagrass, St. Augustinegrass, Bahiagrass and Centipedegrass:** Apply 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse Seasonal conditions. The second application of 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq. ft.

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage. Do not exceed the application rate for any area.

### SPREADER GUIDE - ONE BAG WILL COVER UP TO 5,000 SQUARE FEET

These settings can be used only as a guide, they are not absolute. The speed at which the applicator walks, spreader condition and the texture of the turf will alter the rate of application. If product remains in spreader after application to recommended square foot coverage, spread remainder over same area going at right angles.

Spreader	Spreader Settings
	Lbs. of product per 1,000 sq. ft.
	4 lbs.
Scotts®/Republic Accugreen (Drop)	4 1/4
Scotts®/Republic Speedy Green (Broadcast)	3 3/4
ferti•lome®/Earthway Ev-N-Spree (Broadcast)	14

The above spreader settings are approximate. Calibrate your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

### BROADLEAF WEEDS

Annual Yellow Sweetclover	Buttonweed	Fiddleneck	Narrowleaf Plantain	Russian Pigweed	Virginia Buttonweed
Aster	Canada Thistle	Florida Pusley	Narrowleaf Vetch	Russian Thistle	Virginia Creeper
Austrian Fieldcress	Carolina Geranium	Frenchweed	Nettle	Scarlet Pimpernel	Virginia Pepperweed
Bedstraw	Carpetweed	Galinsoga	Orange Hawkweed	Scotch Thistle	Wavyleaf Bulbthistle
Beggaricks	Catchweed Bedstraw	Goldenrod	Oriental Cocklebur	Sheep Sorrel	Sheeperspurse
Betony, Florida	Catnip	Ground Ivy	Oxalis	Slender Plantain	Western Clematis
Bindweed, Field	Catsear	Gumweed	Parsley-Piert	Smallflower Galinsoga	Western Salsify
Bird Vetch	Chickweed, Common	Hairy Fleabane	Parsnip	Smooth Dock	White Mustard
Bitter Wintercress	Chickweed, Mouseear	Hawkweed	Pearwort	Smooth Pigweed	Wild Aster
Bittercress, Hairy	Chickweed, Mouseear	Healall	Pennycress	Sorrel	Wild Buckwheat
Bitterweed	Chicory	Heartleaf Drymary	Pepperweed	Sowthistle	Wild Carrot
Black-Eyed Susan	Cinquefoil	Heath Aster	Pigweed	Spanishneedles	Wild Four-O'Clock
Black Medic	Clover, Crimson	Hedge Bindweed	Pineywoods Bedstraw	Speedwell	Wild Garlic
Black Mustard	Clover, Hop	Hedge Mustard	Speedwell	Spiny Amaranth	Wild Geranium
Blackseed Plantain	Clover, Red	Henbit	Plains Coreopsis	Spiny Cocklebur	Wild Lettuce
Blessed Thistle	Clover, Strawberry	Hoary Cross	Plantain	Spiny Sowthistle	Wild Marigold
Bloodflower Milkweed	Clover, Sweet	Hoary Plantain	Poison Ivy	Poison Oak	Wild Mustard
Blue Lettuce	Clover, White	Hoary Vervain	Poison Oak	Pokeweed	Wild Onion
Blue Vervain	Cocklebur	Horsenettle	Pokeweed	Spotted Spurge	Wild Parsnip
Bracted Plantain	Common Mullein	Jimsonweed	Poorjoe	Spurweed	Wild Radish
Bristly Oxtongue	Creeping Jenny	Knawel	Prairie Sunflower	Stinging Nettle	Wild Rape
Broadleaf Dock	Cudweed	Knotweed	Picky Lettuce	Strawberry, India Mock	Wild Strawberry
Broadleaf Plantain	Curly Dock	Kochia	Prickly Sida	Tall Nettle	Wild Sweet Potato
Broomweed	Daisy, English	Lambsquarters	Prostrate	Tall Vervain	Wild Vetch
Buckhorn	Daisy, Fleabane	Lespedeza	Prostrate Pigweed	Tansy Ragwort	Woodsorrel
Buckhorn Plantain	Daisy, Oxeye	Mallow	Prostrate Spurge	Tansy Mustard	Woolly Croton
Bulbous Buttercup	Dandelion	Matchweed	Prostrate Vervain	Tanweed	Woolly Morningglory
Bull Thistle	Dichondra	Mexicanweed	Puncturevine	Thistle	Woolly Plantain
Bullnettle	Dogfennel	Milk Vetch	Purslane, Common	Trailing Crownvetch	Wormseed
Burclover	Dollarweed	Morningglory	Ragweed	Tumble Mustard	Yarrow
Burdock	False Dandelion	Mouseear Hawkweed	Red Sorrel	Tumble Pigweed	Yellow Rocket
Burning Nettle	Falseflax	Mugwort	Redroot Pigweed	Velvetleaf	Yellowflower Pepperweed
Burweed	False Sunflower	Musk Thistle	Redstem Filaree	Venice Mallow	
Buttercup		Mustard	Rough Cinquefoil		
			Rough Fleabane		

## 25-0-4

GUARANTEED ANALYSIS

Total Nitrogen (N).....	25%
2% Ammoniacal Nitrogen	
23% Urea Nitrogen*	
Soluble Potash (K <sub>2</sub> O).....	4%
Boron (B).....	0.02%
Copper (Cu).....	0.05%
0.05% Water Soluble Copper (Cu)	
Iron (Fe).....	0.10%
0.10% Water Soluble Iron (Fe)	
Manganese (Mn).....	0.05%
0.05% Water Soluble Manganese (Mn)	
Zinc (Zn).....	0.05%
0.05% Water Soluble Zinc (Zn)	

Derived from: Ammonium Sulfate, Urea, Polymer Coated Sulfur Coated Urea, Muriate of Potash, Sodium Borate, Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, Zinc Sulfate.  
\*7.5% Slowly Available Nitrogen derived from Polymer Coated Sulfur Coated Urea.

F370

### LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price paid will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

ferti•lome® is a registered trademark of Voluntary Purchasing Groups, Inc.

10921-0515-TP



Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418

Visit Us At: [www.fertilome.com](http://www.fertilome.com)

EPA Reg. No. 2217-603-7401 EPA Est. No. 7401-TX-01

Product Questions? 855-270-4776

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapco.org/metals.htm>

10921